




Hypersensitivity risks in biopharmaceutical development

All proteins are potentially immunogenic

The growing number of biological drugs in clinical development is presenting new testing challenges for their manufacturers. As all proteins are potentially immunogenic, the risk that they may induce an unwanted product-related immune response is ever present. The most severe immune response, acute anaphylactic reaction, may be fatal to the patient. What's more, patient and disease-related factors plus process-related impurities also increase the risk of a serious adverse reaction.

Need for better monitoring

The potential severity of the medical consequences, plus the biological complexity of the immunogenic risk, has prompted regulatory authorities to act. They now issue recommendations for performing a systematic immunogenicity assessment from a marketing authorization perspective. Assessing hypersensitivity is thus an increasingly important factor when seeking regulatory approval for a new biologic.



Key questions about the risks of hypersensitivity

Developers of new biologics need to address a number of important questions relating to hypersensitivity and the risk factors that accompany it.

These questions include:

- **How do I know that this protein is not allergenic?**
- **How do I test for allergenicity during a clinical trial?**
- **How do I ensure that this drug will not cause allergic reactions post-market?**


Great performance and risk management go hand in hand

Sensitive *in vitro* IgE tests will assess IgE levels before allergenic reactions occur, alerting developers to potential risks and thereby helping manage them. Phadia is the world-leader in allergy testing with a 75 % market share. Our allergy tests are the most sensitive on the market and our experience is unrivalled. With more than 40 years of R&D in allergy pathology and assay development combined with manufacturing and a complete supply chain, Phadia is your ideal partner during drug development.

Challenges

- Must distinguish between an infusion reaction and anaphylaxis due to the very different clinical consequences
- Pre-sensitization caused by previous exposure to similar or related proteins
- Therapeutic proteins used for long-term, intermittent treatment
- Therapeutic proteins used for substitution of an endogenous protein
- Product-related factors influence immunogenicity
- Process-related impurities present





Your partner in allergy risk management and immunogenicity testing

Phadia

Phadia AB, P O Box 6460, SE-751 37 Uppsala, Sweden
Tel +46 18 16 50 00. www.phadia.com

Head office Sweden +46 18 16 50 00 Austria +43 1 270 20 20 Belgium +32 2 749 55 15 Brazil +55 11 3345 5050 Czech Republic +420 220 511 392 Denmark +45 7023 3306 Finland +358 9 8520 2560 France +33 1 61 37 34 30 Germany +49 761 47 8050 Great Britain/Ireland +44 1 908 769 110 Italy +39 02 641 634 11 Japan +81 3 5365 8332 Korea +82 2 2027 5400 Netherlands +31 30 602 3700 Norway +47 21 67 32 80 Portugal +351 21 423 5350 South Africa +27 11 793 5337 Spain +34 935 765 800 Sweden +46 18 16 50 00 Switzerland +41 43 343 4050 Taiwan +886 2 2516 0925 United States +1 800 346 4364 Other +46 18 16 56 16

ImmunoCAP is a registered trademark of Phadia AB. EIIA is a trademark of Phadia AB.